

Risk Categories for COVID-19		
	Exposure to a symptomatic confirmed case	Management
High	Living in the same household as, being an intimate partner of, or providing care in a non-healthcare setting (such as a home) for a person with symptomatic laboratory-confirmed COVID-19 infection <i>without using recommended precautions</i> for <a href="#">home care</a> and <a href="#">home isolation</a>	<p><b>Active Monitoring - Daily contact with the health department</b></p> <p>Restriction to home</p> <p>Immediate isolation and notification to the health department if symptomatic.</p>
Medium	Close contact with a person with symptomatic laboratory-confirmed COVID-19. Living in the same household as, an intimate partner of, or caring for a person in a non-healthcare setting (such as a home) to a person with symptomatic laboratory-confirmed COVID-19 infection <i>while consistently using recommended precautions</i> for <a href="#">home care</a> and <a href="#">home isolation</a>	<p><b>Self-Monitoring- take temperature daily watch for signs and symptoms</b></p> <p>Recommendation to remain at home or in a comparable setting</p> <p>Practice social distancing</p> <p>Immediate isolation and notification to the health department if symptomatic.</p>
Low	Being in the same indoor environment (e.g., a classroom, a hospital waiting room) as a person with symptomatic laboratory-confirmed COVID-19 for a prolonged period of time but not meeting the definition of close contact	<p><b>Self-observation self-assess for temperature and symptoms</b></p> <p>Practice social distancing</p> <p>Immediate isolation if becomes symptomatic. Stay at home. Seek medical advice if you develop severe symptoms.</p>